

LA-UR-18-23274

Approved for public release; distribution is unlimited.

Title: Stand Up! and move Why we need to get up out of our chairs ...and move

Author(s): Hite, Peter Wesley

Intended for: Research project was made as part of my master's course work. It was

presented to ADBI WSST on 4/12. My professor will review and grade.

Issued: 2018-04-17



Stand Up! and move

Why we need to get up out of our chairs

...and move



MY PATIENT'S WELL-BEING IS MY FIRST PRIORITY.

I PROVIDE THOUGHTFUL, COMPASSIONATE HEALTH CARE, ALWAYS RESPECTING THE RIGHTS, WELFARE & DIGNITY OF OTHERS.



AS THE ADVOCATE FOR MY PATIENT'S BEST MEDICAL INTEREST, I MAKE COMPETENT DECISIONS BASED ON EVIDENCE-BASED PRACTICE.

I ACT WITH INTEGRITY.

I FULLY UNDERSTAND AND UPHOLD THE NATA CODE OF ETHICS, PROVIDING

THE BEST POSSIBLE PATIENT CARE AT ALL TIMES.

I COMPLY WITH THE LAWS AND REGULATIONS GOVERNING THE PRACTICE OF ATHLETIC TRAINING, AND I PLEDGE TO MAINTAIN AND PROMOTE THE HIGHEST QUALITY OF HEALTH CARE.







UNCLASSIFIED





Stand Up! and move

- Sedentary behavior is defined as any waking behavior characterized by an energy expenditure of 1.5 METS or less while in a sitting or reclining posture¹
- Sitting and typing uses 1.45 METS²
- Standing Still 1.59 METS²
- Walking 1MPH 2.66 METS²
- Climbing stairs 4 METS¹⁴
- Running (8MPH) 13.5¹⁴






I'm just sitting down, how is that risky?

- 60% of computer users reported cumulative trauma disorders (ie musculoskeletal disorders and eye soreness)³
- 33% of all worker injury/illness cases caused by MSD⁴
- Increased risk of acute or transient pain complaints when keying or using mouse intensively⁵
- Sedentary behavior brings high risk of cardiovascular disease, stroke, colon cancer, diabetes (300,000 deaths annually linked to physical inactivity)⁶
- Decrease in mental wellbeing linked to physical inactivity⁷





How Do These Things Become Harmful?





How do Musculoskeletal Disorders ("ergonomic injuries") Develop?



- Trauma From Work Accumulates in Tissue
 - Poor work design increases build-up
- Tissue Can "Hold" a Limited Amount of Trauma Before Injury
 - Fitness contributes a small increase in trauma capacity



What can I do?

- Limit sitting time to a max of 2 hours, stand up and move after 30 minutes of uninterrupted sitting⁸
- Take time to stretch when you're at rest
- During your break take a brief walk, use the stairs if you can
- Microbreaks can benefit health and mental productivity, and reduce eye strain^{9,10,11}
- Sit-to-Stand Desks have shown a reduction in physical discomfort compared to sitting and a reduction in spinal compressive forces ^{12,13}










LANL COMBINED RECORDABLE SOURCE ANALYSIS FEBRUARY 2017 - JANUARY 2018





UNCLASSIFIED




Tools

- Apps: Stand Up! iOS App Apple app store
- Apps: Twenty: Stand up every 20 mins Android AppGoogle Play
- Software: Stretch Break
- Websites: e.ggtimer.com
- Wellness center
- Ergo eval/ergo equipment











UNCLASSIFIED

For more info:

Pete Hite, LAT, ATC

ADBI Ergo Guy 505-551-5368

PeteHite@lanl.gov

ergo.lanl.gov





UNCLASSIFIED

References

- 1. Tremblay, M. (2012). Letter to the Editor: Standardized use of the terms "sedentary" and "sedentary behaviours". *Applied Physiology, Nutrition, and Metabolism*, 37(3), pp.540-542.
- Mansoubi, M., Pearson, N., Clemes, S., Biddle, S., Bodicoat, D., Tolfrey, K., Edwardson, C. and Yates, T. (2018). *Energy expenditure during common sitting and standing tasks: examining the 1.5 MET definition of sedentary behaviour.* [online] BCM Public Health. Available at: https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-015-1851-x
- Institute of Occupational Safety and Health. *The study of fatigue and physiological conditions of VDT operators* Institute of Occupational Safety and Health, Taipei (1996)
- Osha.gov. (2018). Safety and Health Topics | Ergonomics | Occupational Safety and Health Administration. [online] Available at: https://www.osha.gov/SLTC/ergonomics/ [Accessed 11 Apr. 2018].
- Andersen, Johan H., et al. "Risk Factors for Neck and Upper Extremity Disorders among Computers Users and the Effect of Interventions: An Overview of Systematic Reviews." PLoS ONE, vol. 6, no. 5, 2011, p. e19691. Health Reference Center Academic, http://link.galegroup.com.prx-usa.lirn.net/apps/doc/A476891676/HRCA?u=lirn55718&sid=HRCA&xid=d8b95e01.
- 6. Cdc.gov. (2018). Physical Inactivity | Gateway to Health Communication | CDC. [online] Available at: https://www.cdc.gov/healthcommunication/toolstemplates/entertainmented/tips/PhysicalInactivity.html
- Harris, M. (2018). The relationship between physical inactivity and mental wellbeing: Findings from a gamification-based community-wide physical activity intervention. *Health Psychology Open*, 5(1), p.205510291775385.
- Plotnikoff, R. and Karunamuni, N. (2012). Reducing Sitting Time: The New Workplace Health Priority. *Archives of Environmental & Occupational Health*, 67(3), pp.125-127.
- 9. Ariga, A. and Lleras, A. (2011). Brief and rare mental "breaks" keep you focused: Deactivation and reactivation of task goals preempt vigilance decrements. *Cognition*, 118(3), pp.439-443.
- BALCI, R. and AGHAZADEH, F. (2003). The effect of work-rest schedules and type of task on the discomfort and performance of VDT users. *Ergonomics*, 46(5), pp.455-465.



References cont.

- Idowu, P. A., Adedoyin, R. A., & Adagunodo, R. E. (2005). Computer-related repetitive strain injuries. *Journal of the Nigeria Society of Physiotherapy*, *15*(1), 13.
- Grunseit, A., Yuk-Yin Chau, J., van der Ploeg, H. and Bauman, A. (2013). "Thinking on your feet": A qualitative evaluation of sit-stand desks in an Australian workplace. *BMC Public Health*, 13(1).
- Callaghan, J. and McGill, S. (2001). Low back joint loading and kinematics during standing and unsupported sitting. *Ergonomics*, 44(3), pp.280-294.
- Harvard Health Publishing; Harvard Medical School(2009). *MET-hour equivalents of various physical activities Harvard Health*. [online] Harvard Health. Available at: https://www.health.harvard.edu/staying-healthy/met-hour-equivalents-of-various-physical-activities [Accessed 12 Apr. 2018].



